

Health Protocol

1. All calves must be weaned a minimum of 30 days or more 45 days is preferred.
2. Calves must weigh a minimum of 550 pounds upon delivery.
3. All bulls must be castrated.
4. All calves must be treated for internal and external parasites.
5. All cattle must be dehorned.
6. Cattle should receive two rounds of modified live viral vaccine given preferably preweaning and at weaning.

Modified Live Virus

- IBR-(Infectious Bovine Rhinotracheitis)
- BVD-(Bovine Viral Diarrhea Virus 2 types)
- P13-(Parainfluenza - not an important pathogen, but always in virals)
- BRSV-(Bovine Respiratory Syncytial Virus)

Examples of Pasteurella vaccine are as follows: Express5, Bovishield Gold 5, Pyramid 5, Vista 5, etc.

****Please consult your veterinarian before using MLV vaccine on calves nursing pregnant cows****

7. Pasteurella hemolytica (Mannheimia hemolytica) and Pasteurella multocida bacterin toxoid are optional. Pasteurella vaccine should only be administered preweaning if used.

Examples of Pasteurella vaccine are as follows: Palmoguard PHM-1, One Shot, etc.

8. 7-way blackleg with histophilus (2 doses required)

Examples of Blackleg vaccine are as follows: Ultrabac7somubac, Varvac7somnus, or use hemophilus separate or in conjunction with virals (Express5HS), etc.

Adjunct procedures/Biologicals **Not required for entry**

****Please consult your veterinarian for recommendations on which products/procedures best fit your livestock operation****